

Washington University in St. Louis

Washington University Open Scholarship

Volume 12

Washington University
Undergraduate Research Digest

Spring 2017

Investigating the Role of $\gamma 3$, and Auxiliary Subunit of the BK Channel, in Myometrial Smooth Muscle Cells

Will Freeman

Washington University in St. Louis

Follow this and additional works at: https://openscholarship.wustl.edu/wuurd_vol12

Recommended Citation

Freeman, Will, "Investigating the Role of $\gamma 3$, and Auxiliary Subunit of the BK Channel, in Myometrial Smooth Muscle Cells" (2017). *Volume 12*. 58.

https://openscholarship.wustl.edu/wuurd_vol12/58

This Abstracts A-I is brought to you for free and open access by the Washington University Undergraduate Research Digest at Washington University Open Scholarship. It has been accepted for inclusion in Volume 12 by an authorized administrator of Washington University Open Scholarship. For more information, please contact digital@wumail.wustl.edu.

INVESTIGATING THE ROLE OF $\gamma 3$, AN AUXILIARY SUBUNIT OF THE BK CHANNEL, IN MYOMETRIAL SMOOTH MUSCLE CELLS

Will Freeman

Mentor: Sarah K. England

The large-conductance Ca^{2+} -activated K^+ (BK) channel is known to be important in regulating contractility and relaxation in myometrial smooth muscle cells (MSMCs) by controlling the flux of K^+ ions across the cell membrane. The BK channel is activated by voltage and intracellular Ca^{2+} concentration ($[\text{Ca}^{2+}]_{\text{in}}$). In addition, the BK channel has been shown to be constitutively active in MSMCs isolated from pregnant women during labor as compared to MSMCs isolated from non-laboring tissue. This underlying mechanism for this constitutive activity in BK activity has not been fully investigated. The BK channel can be modulated by auxiliary proteins, of which one type, the γ subunits, have been shown to activate the BK channel at relatively hyperpolarized membrane potentials and at very low $[\text{Ca}^{2+}]_{\text{in}}$. Here we studied whether association with the γ subunits could underlie the BK channel phenotype seen at labor. We focused on the $\gamma 3$ subunit since it was reported to have the highest abundance in uterine smooth muscle. We investigated the expression and role of the $\gamma 3$ subunit in non-laboring (NL) MSMCs using both molecular biology and electrophysiological techniques. Our first experiments assessed whether $\gamma 3$ protein was present in hMSMCs, however we were unsuccessful due to the non-specificity of the antibodies targeted against $\gamma 3$. To identify the functional role of the $\gamma 3$ subunit, BK channel currents were measured using patch-clamp in the inside-out excised-patch configuration. We recorded currents from HEK cells overexpressed with BK channel alone and the BK channel co-expressed with the $\gamma 3$ subunit, but did not detect a change in BK activity in the presence of the $\gamma 3$ subunit. We also investigated whether the channel activity changed with a 1:4 ratio and from NL MSMC. Future studies will determine whether the association of the $\gamma 3$ is present in laboring tissue.